Table 6-5.1. Fertilizer Requirements

TYPE OF SPECIES	YEAR	ANALYSIS OR EQUIVALENT N-P-K	RATE	N TOP DRESSING RATE
Cool season grasses	First Second Maintenance	6-12-12 6-12-12 10-10-10	1500 lbs./ac. 1000 lbs./ac. 400 lbs./ac.	50-100 lbs./ac. 1/2/ 30
Cool season grasses and legumes	First Second Maintenance	6-12-12 0-10-10 0-10-10	1500 lbs./ac. 1000 lbs./ac. 400 lbs./ac.	0-50 lbs./ac. 1/ — —
3. Ground covers	First Second Maintenance	10-10-10 10-10-10 10-10-10	1300 lbs./ac. 3/ 1300 lbs./ac. 3/ 1100 lbs./ac.	_ _ _
4. Pine seedlings	First	20-10-5	one 21-gram pellet per seedling placed in the closing hole	_
5. Shrub Lespedeza	First Maintenance	0-10-10 0-10-10	700 lbs./ac. 700 lbs./ac. 4/	_
Temporary cover crops seeded alone	First	10-10-10	500 lbs./ac.	30 lbs./ac. 5/
7. Warm season grasses	First Second Maintenance	6-12-12 6-12-12 10-10-10	1500 lbs./ac. 800 lbs./ac. 400 lbs./ac.	50-100 lbs./ac. 2/6/ 50-100 lbs./ac. 2/ 30 lbs./ac.
Warm season grasses and legumes	First Second Maintenance	6-12-12 0-10-10 0-10-10	1500 lbs./ac. 1000 lbs./ac. 400 lbs./ac.	50 lbs./ac./6/

^{1/} Apply in spring following seeding.

^{2/} Apply in split applications when high rates are used.

^{3/} Apply in 3 split applications.

^{4/} Apply when plants are pruned.

^{5/} Apply to grass species only.

^{6/} Apply when plants grow to a height of 2 to 4 inches.

Table 6-5.2 - Permanent Cover

PLANT, PLANTING RATES, AND PLANTING DATES FOR PERMANENT COVER

	<u>1/ - F</u>	PLS 2/		Plan	ing D	ates	by R	esou	ce A	reas	Plant	ting [Dates		Τ	
Species	Per Acre	Per 1000 sq.ft.	(Sc	olid lir	nes in		e opti						indic	ate		<u>Remarks</u>
BAHIA, PENSACOLA		<u>34.11.</u>	J	F					J				N	D	+	
(Paspalum notatum)						, i				-						
alone or with temporary cover	60 lbs.	1.4 lb.														166,000 seed per pound. Low growing. Sod forming. Slow to establish. Plant with a companion crop. Will spread into bermuda pastures and lawns. Mix with Sericea lespedeza or weeping lovegrass.
with other perenials	30 lbs.	0.7 lb.													1	
BAHIA, WILMINGTON			J	F	М	Α	М	J	J	Α	S	0	N	D		
(Paspalum notatum)															1	166,000 seed per pound. Low growing. Sod forming. Slow to establish. Plant with a companion crop. Will
alone or with temporary cover	60 lbs.	1.4 lb.														spread into bermuda pastures and lawns. Mix with Sericea lespedeza or weeping lovegrass.
with other perenials	30 lbs.	0.7 lb.														
BERMUDA, COMMON			J	F	М	Α	М	٦	J	Α	S	0	N	D		
(Cynodon dactylon)																
Hulled seed															1	1,787,000 seed per pound. Quick cover. Low growing and sod forming. Full sun. Good for athletic fields.
	10 lbs.	0.2 lb.														
with other perenials	6 lbs.	0.1 lb.														
BERMUDA, COMMON			J	F	М	Α	М	٦	J	Α	S	0	N	D		
(Cynodon dactylon)																
Unhulled seed															F	Plant with winter annuals. Plant with tall fescue.
with temporary cover	10 lbs.	0.2 lb.														
with other perenials	6 lbs.	0.1 lb.														
BERMUDA, SPRIGS			J	F	М	Α	М	J	J	Α	S	0	Ν	D		
(Cynodon dactylon)	40 cu. ft.	0.9 cu.ft.													,	A cubic foot contains approximately 650 sprigs. A bushel contains 1.25 cubic feet or approximately 500 sprigs.
Coastal, Common, or Tift 44		or													A	A cubic foot contains approximately 650 sprigs. A businer contains 1.25 cubic feet of approximately 500 sprigs.
Coastal, Common, or Tilt 44	sod plu	ıgs 3' X 3'														
CENTIPEDE			J	F	М	Α	М	J	J	Α	S	0	Ν	D		Drought tolerant. Full sun or partial shade. Effective adjacent to concrete and in concentrated flow areas.
(Eremochloa ophiuroides)	Block	sod only														Irrigation is needed until fully estabilzhed. Do not plant near pastures. Winterhardy as far north as Athens and Atlanta.
CROWNVETCH			J	F	М	Α	М	J	J	Α	S	0	N	D		лини.
(Coronilla varia)				<u> </u>				Ĭ	Ť		Ĭ	Ü	Ë	Ť	1	
with winter																
annuals or cool	15 lbs/	0.3 lb.														100,000 seed per pounf. Dense growth. Drought tolerant and fire resistant. Attractive rose, pink, and white
season grasses																blossoms spring to late fall. Mix with 30 lbs of Tall Fescue or 15 lbs of rye. Innoculate seed with M inoculant. Use from North Atlanta and Northward.
																030 HOIH NOITH Atlanta and NOITHWAID.
															\perp	

PLANT, PLANTING RATES, AND PLANTING DATES FOR PERMANENT COVER

	1/	- PLS 2/		Plan	ting D	ates	by R	esou	rce A	reas	Plan	ting	Dates	<u> </u>	
Species	_	re Per 1000	(Sc	olid lir	nes in								indic	ate	<u>Remarks</u>
		sq.ft.						e but							
FESCUE, TALL			J	F	М	Α	М	J	J	Α	S	0	Ν	D	
(Festuca arundinacea)															
alone	50 lbs.	1.1 lbs.													
	00 11	0.7.11											-		
with other perennials	30 lbs.	0.7 lb.													227,000 seed per pound. Use alone only on better sites. Not for drought soils. Mix with perennial lespedezas or crownvetch. Apply topdressing in spring following fall plantings. Not for heavy use areas or athletic fields.
LESPEDEZA, SERICEA			J	F	М	Α	М	J	J	Α	S	0	N	D	
(Lespedeza cuneata)															
scarified	60 lbs	. 1.4 lbs.													350,000 seed per pound. Widely adapted. Low maintenance. Mix with weeping lovegrass, common bermuda, bahia, or tall fescue. Takes 2 - 3 years to become fully established. Excellent on road banks. Inoculate seed with EL inoculant.
unscarified	75 lbs	. 1.7 lb.													Mix with Tall fescue or winter annuals.
seed-bearing hay	3 tons	138 lbs.			-										Cut when seed is mature, but before it shatters. Add Tall fescue to winter annuals.
LESPEDEZA			J	F	М	Α	М	J	J	Α	S	0	N	D	
Ambro virgata															
(Ledpedeza virgata DC)															
or															
Appalow (Lespedza cuneata															300,000 seed per pound. Height of growth is 18 - 24 inches. Advantageous in urban areas. Spreading-type
[Dumont] G. Don)															growth has bronze coloration. Mix with Weeping lovegrass, Common bermuda, bahia, tall fescue, or winter
,															annuals. Do not mix with Sericea lespedaza. Slow to develop solid stands. Inoculate seed with EL inoculate.
scarified	60 lbs	. 1.4 lb.													
unscarified	75 lbs	. 1.7 lb.													

PLANT, PLANTING RATES, AND PLANTING DATES FOR PERMANENT COVER

	1/ - P	LS 2/	1	Plan	ing Da	ates	by R	esou	rce A	reas	Plan	ting [Dates		
Species	Per Acre		(Sc	olid lir	nes inc								indic	ate	<u>Remarks</u>
		sq.ft.		permissbale but marginal dates.)											
LESPEDEZA, SHRUB			J	F	М	Α	М	J	J	Α	S	0	Ν	D	
(Lespedeza bicolor)															
(Lespedeza thumbergii)															Provide wildlife food and cover.
plants	3' >	x 3'													
LOVEGRASS, WEEPING			J	F	М	Α	М	J	7	Α	S	0	Ν	D	
(Eragrostis curvula)															
alone	4 lbs.	0.1 lb.													1,500,000 seed per pound. Quick cover. Drought tolerant. Grows well with Sericea lespedeza on road banks.
with other perennials	2 lbs.	0.05 lb.													
MAIDENCANE			J	F	М	Α	М	J	J	Α	S	0	Ν	D	
(Panicum hemitomon)															For very wet sites. May clog channels. Dig sprigs from local sources. Use along river banks and shorelines.
sprigs	2' x 3' s	spacing													
PANICGRASS,			J	F	М	Α	М	٦	7	Α	S	0	Ν	D	
ATLANTIC COASTAL															Grows well on coastal sand dunes, borrow areas, and gravel pits. Provides winter cover for wildlife. Mix with
(Panicum amarum var.	20 lbs.	0.5 lb.													Sericea lespedeza except on sand dunes.
amarulum)	20 1301	0.0 .0.													
REED CANARY GRASS	1		J	F	М	Α	М	J	J	Α	S	0	N	D	
(Phalaris arundinacea)															
alone	50 lbs.	1.1 lb.													Grows similar to tall fescue.
with other perennials	30 lbs.	0.7 lb.													
SUNFLOWER, 'AZTEC'			J	F	М	Α	М	J	J	Α	S	0	N	D	
MAXIMILLIAM	10 lbs.	0.2 lb.													227,000 seed per pound. Mix with weeping lovegrass or other low-growing grasses or legumes.
(Helianthus maximiliani)															

Table 6-4.1 - Temporary Cover for Companion Crops¹

PLANT, PLANTING RATES, AND PLANTING DATES FOR TEMPORARY COVER OR COMPANION CROPS¹

Species	Broadcas PL			Plan	ting [Dates	by R	esou	rce A	reas	Plan	ting E	ates		Remarks				
3,73.00	Per Acre	Per 1000 sq.ft.	(S	olid liı		ndicat permis									i terraino				
MILLET, BROWNTOP			J	F	M	Α	M	J	J	Α	S	0	Ν	D					
(Panicum fasciculatum)																			
alone	40 lbs.	0.9 lb.													137,000 seed per pound. Quick dense cover. Will provide too much competition in mixtures if seeded at high rates.				
in mixtures	10 lbs.	0.2 lb.																	
MILLET, PEARL			J	F	М	Α	М	J	J	Α	S	0	Ν						
(Pennesetum glaucum)															88,000 seed per pound. Quick dense cover. May reach 5 feet in height. Not recommended for mixtures.				
alone	50 lbs	1.1 lb.																	
OATS			J	F	M	Α	M	J	J	Α	S	0	Ν	D					
(Avena sativa)																			
alone	4 bu. (128 lbs.)	2.9 lb.													13,000 seed per pound. Use on productive soils. Not as winter hardy as rye or barley.				
in mixtures	1 bu. (32 lbs.)	0.7 lb.																	
RYE			J	F	М	Α	M	J	J	Α	S	0	N	D					
(Secale cereale)																			
alone	3 bu. 168 (lbs.)	3.9													18,000 seed per pound. Quick cover. Drought tolerant and winter hardy.				
in mixtures	1/2 bu. 28 lbs.	0.6 lb.																	
RYEGRASS, ANNUAL			J	F	M	Α	М	٦	J	Α	S	0	Ν	D					
(Lolium temulentum)		С																	
alone	40 lbs.	0.9 lb													227,000 seed per pound. Dense cover. Very competitive and is not to be used in mixtures.				
SUDANGRASS			J	F	М	Α	М	J	J	Α	S	0	N	D					
(Sorghum Sudanese)															55,000 seed per pound. Good on drought sites. Not recommended for mixtures.				
alone	60 lbs.	1.4 lb.									İ								
WHEAT			J	F	М	Α	М	J	J	Α	S	0	N	D					
(Triticum Aestivum)																			
alone	3bu. (180 lbs.)	4.1 lb													15,000 seed per pound.				
in mixtures	1/2 bu. (30 lbs.)	0.7 lb.																	

¹ Temporary cover crops are very competitive and will crowd our perennials if seeded too heavily

² Reduce seeding rates by 50% when drilled.

³ PLS is an abbreviation for Pure Live Seed.